

Regional Climate Projections over Spain: Atmosphere. Future Climate Projections

Jesús Fernández¹, Ana Casanueva^{1*}, Juan Pedro Montávez², Miguel Ángel Gaertner³, María Jesús Casado⁴, Rodrigo Manzanas⁵, José Manuel Gutiérrez⁵

¹ Meteorology Group. Dept. Applied Mathematics and Comp. Sci. Universidad de Cantabria, Spain

² Regional Atmospheric Modelling Group, Department of Physics, University of Murcia, Spain

³ Faculty of Environmental Sciences and Biochemistry. University of Castilla-La Mancha, Spain

⁴ Agencia Estatal de Meteorología (AEMET), Spain

⁵ Meteorology Group. Instituto de Física de Cantabria (CSIC-UC), Spain

(*) presently at the Federal Office of Meteorology and Climatology MeteoSwiss, Switzerland

Introduction

Regional climate information is increasingly being demanded by vulnerability, impact and adaptation (VIA) research communities. This information is required to feed impact models for specific sectors (health, energy, food availability, risk management, water resources) and for decision-making processes at different levels. Different global and regional climate change projections for the 21st century have been produced over the last decades using both dynamical regional climate models (RCMs) and statistical downscaling methods (SDMs) in a series of international and national initiatives. As a result, large ensembles of future global (e.g. CMIP3, CMIP5) and regional (e.g. ENSEMBLES, CORDEX) climate projections are available, sampling most of the uncertainties affecting climate change. Nowadays, users are confronted with the technical and ethical (distillation) dilemma of deciding which information out of the large amount of available data is best suited for their application, in order to address the different sources of uncertainty for their specific region/problem (Hewitson, 2013).

Regional Climate Change Scenarios for Spain

The Earth System Grid Federation (ESGF, <https://esgf.llnl.gov>) infrastructure provides standardized access to the last-generation climate change model output data (from CMIP5 and CORDEX). However, the direct use of ESGF is still complex and slow for an average user and therefore there are still several portals providing scenario data (e.g. for a particular region), as well as sector-specific derived indices (Hewitson et al., 2017).

In Spain, the main sources of regional climate change information have been collected by the Escenarios-PNACC initiative. The first dataset (Escenarios-PNACC, 2012) was based on CMIP3 projections from global circulation models (using B1, A1B and A2 greenhouse gases emission scenarios), and provided information

for temperature and precipitation and several derived indices (e.g. percentiles). Besides the European-wide dynamical regional climate change projections provided by the EU-funded ENSEMBLES project (Déqué et al., 2012), and the contribution from AEMET (statistical downscaling methods), this initiative was fed with the outcomes of two strategic actions (ESCENA and ESTCENA, for dynamical and statistical downscaling, respectively) undertaken by the Spanish scientific community on regional climate change projection in the framework of the Spanish R+D 2008–2011 Program. The dynamical simulations complemented those produced in ENSEMBLES, focusing on a domain centered in the Iberian Peninsula (Fernández et al. 2012; Jiménez-Guerrero et al. 2013, Domínguez et al., 2013). The statistical downscaling scenarios were produced with different analog- and regression-based downscaling methods (Gutiérrez et al. 2012). The observational data set was based on a selection of stations provided by AEMET and on the gridded observations provided by Spain02 v2.1 (regular grids, at a 20 km resolution; see Herrera et al. 2012). More information and links to access data are described in Escenarios-PNACC (2012).

There have been a number of studies analysing these results and assessing the limitations of the different datasets forming Escenarios-PNACC 2012, focusing mainly on temperature and precipitation. For instance, Turco et al. (2015) analysed the ENSEMBLES regional projections for daily maximum temperature and precipitation over Spain, and found consistent changes up to 2050 (A1B scenario) among the different members, generally indicating a decrease in precipitation (between -5 and -25 %) and an increase in maximum temperature (between 1 and 2.5°C, depending on the season/area). Gutiérrez et al. (2013) reported a limitation of generic analog based methods to extrapolate future temperatures for the last decades of the century. Therefore, those

There have been also some studies analysing additional variables such as snow, wind speed or dry spells. For instance, Pons et al. (2016) reported a decreasing trend in annual snowfall frequency (measured as the annual number of snowfall days) from the ENSEMBLES regional projections, with member values ranging from -3.7 to -0.5 days decade⁻¹ (-2.0 day decade⁻¹ for the ensemble mean). These future trends are similar to the historical observed ones since 1970 (-2.2 days decade⁻¹), and are mainly determined by the increasing temperatures. Gómez et al. (2016) analysed wind speed using ENSEMBLES data and found that the wind speed for 2031–2050 is reduced up to 5% compared to the 1980–1999 control period for all models. The models also agree on the time evolution of spatially averaged wind speed in each region, showing a negative trend for all of them. López-Franca (2015) analysed dry spells over Spain and found an increase of the probability of occurrence of long dry spells together with a decrease in the shorter ones.

These studies provide comprehensive information on the Escenarios-PNACC dataset, although further research is needed to understand the methodological and practical limitations of regional projections, particularly the extrapolation capability of the different methods, i.e. the robustness of the stationarity assumption.

dynamical projections for Spain with the MM5 model. Pérez (2014) and Gonçalves (2014) describe dynamical downscaling results using WRF over the Canary Islands and Mediterranean Spain, respectively. Ramos et al. (2013) analysed CMIP3 precipitation projections with a SDM.

An update of Escenarios-PNACC is being prepared, based on the new information produced using CMIP5 global projections (focused on the RCP2.6, RCP4.5 and RCP8.5 scenarios). The new edition (Escenarios-PNACC 2017) is based on the research done by Spanish groups in the framework of the EU VALUE SDM intercomparison project (Maraun et al., 2015), and the RCM projections produced in the context of the EURO-CORDEX (Jacob et al., 2014) and MED-CORDEX (Ruti et al., 2016) initiatives. The update of the observational dataset is Spain02 v5 (at 10km resolution, both regular and rotated grids). Most of this information is already available (with daily temporal resolution) at http://www.aemet.es/es/serviciosclimaticos/cambio_climat and <http://www.meteo.unican.es/escenarios-pnacc>




Figure 1: GCM-RCM matrix for the dynamical (blue from EURO-CORDEX, green from MED-CORDEX) and statistical (red) ensemble of regional projections from CMIP5 data available as of April 30th, 2017. Numbers show the number of ensemble members (marginal frequencies) for a given GCM (rows) or RCM/SDM (columns). The available projections comprise simulations at three resolutions (0.11°, 0.22° and 0.44°) and for three RCP scenarios (RCP 2.6, 4.5 and 8.5), as displayed in the matrix shown in the upper-left corner. Note that statistical downscaling methods provide local (station-based) information (denoted by the points over the square boxes) and, in some cases, also areal grid-box information (downscaled from Spain02 v5). Additionally, AEMET statistical downscaling point estimates are also available for 14 other CMIP5 GCMs for ANA and 10 for SDSM, not shown in this table.




Figure 2: Projected changes for seasonal temperature ($^{\circ}\text{C}$, top) and precipitation ($\%$, bottom) by 2021- 2050, with respect to the average of the 1971-2000 period, on average for continental Spain and the Balearic Islands for two different forcing scenarios (RCP8.5, RCP4.5 and both "All") considering the GCM, SDM, or RCM projections (in different rows for each scenario). In order to provide comprehensive information about the ensemble uncertainty in the climate change signal, different percentiles (5, 25, 75 and 95) are given for each case. The range 25-75 is the typical value used to characterize the ensemble spread, whereas 5th and 95th values characterize the extreme signals within the ensemble

Figure 1 shows the information already available from Escenarios-PNACC 2017, with a GCM-RCM/SDM matrix for the multi-project ensemble considered in this study and publicly available as of April 30th, 2017. The numbers show the marginal contribution to the ensemble for each particular GCM (in rows) and RCM/SDM (in columns). These include different forcing scenarios and RCM resolutions (see figure caption). For each GCM-RCM/SDM

couple, boxes indicate the available spatial resolutions (0.11° , 0.22° and 0.44°) and RCP scenarios (RCP 2.6, 4.5 and 8.5). Note that, besides the two spatial resolutions, SDMs also provide local (station based) projections for a number of stations (over 2300 selected by AEMET). This makes a total of 255 regional projections available, forming a large multi-model multi-downscaling ensemble sampling the regional uncertainty for climate

change projections over Spain. This dataset will be the basis for future regional climate change studies in Spain.

Figure 2 shows projected temperature and precipitation changes by 2021-2050 averaged over continental Spain and the Balearic Islands. It includes only the set of GCMs downscaled by dynamical and statistical models for the same RCP and spatial resolution, thus forming a balanced ensemble representative of regional uncertainties (grey shadow in Figure 1). Projected changes are presented as central 50% and 90% ranges (obtained from the 25th-75th and 5th-95th percentiles of the ensemble), estimated by different sub-ensembles, and for different seasons. For precipitation, there is a general trend to decreasing average precipitation in all seasons, although the expected range of change is only completely on the drier side during summer, reaching an average decrease of -30% for RCM estimates. Regarding temperature, the largest increases are expected in summer and autumn, reaching close to 3°C with respect to the 1971-2000 climatology in the upper end, and no less than 1°C in the most conservative estimates. GCMs tend to provide warmer estimates, which are matched in the upper end by SDMs. These, however, show larger spread and colder lower ends. RCMs provide, in general, colder estimates.

A description of the EURO-CORDEX projections at European level is given in Jacob et al. (2014), which also presents a comparison of the results with the previous ENSEMBLES scenarios. Some statistical downscaling studies building on CMIP5 data have been recently published analysing different aspects of climate change projections in Spain. For extremes, Monjo et al. (2016) applied a two-step analogue/regression downscaling statistical downscaling using CMIP5 predictors over Spain, and analysed 50 and 100-year return values for precipitation. They found that the projected changes were in general smaller than the natural variability. Future changes in extremes such as tropical-like cyclones phenomena (known as “medicanes” when located over the Mediterranean sea) in oceanic regions around the Iberian Peninsula have been studied with CMIP5 models (Romero and Emanuel, 2017) or ENSEMBLES RCMs (Romera et al., 2017)

Impact Studies and Bias Correction

There have been also a number of studies analysing the impact of climate change projections in different sectors using climate-related indices building on regional climate information. For instance, Bedia et al. (2013, 2014) analysed fire danger projections over Spain using the Fire Weather Index (FWI), and statistical/dynamical projections with CMIP3/ENSEMBLES data, respectively. Resco et al. (2015) and Lorenzo et al. (2016) analysed winegrowing in Spain using different bioclimatic indices and ENSEMBLES data. Jerez et al. (2015) analysed photovoltaic power generation in Europe using EURO-

CORDEX data. Esteve-Selma (2012) studied the future distribution of *Tetraclinis articulata* (an endemic Mediterranean tree). Bafaluy (2014) investigated a number of climate indices relevant for tourism using ENSEMBLES data. Casanueva et al. (2014) introduced the direct application of statistical downscaling to multi-variate climate indices for fire danger and tourism.

One of the main problems found in impact studies when analysing climate-derived indices using RCM data is the effects introduced by model biases. Casanueva et al. (2016) provides a detailed analysis of EURO-CORDEX biases over Spain. This has motivated a huge research activity over the last two decades in order to find suitable bias adjustment methods, able to correct model biases against a reference high-resolution climatology. An intercomparison of different bias adjustment methods over Europe, using ENSEMBLES data, is provided in Dosio et al. (2012). Several impact studies have explored the use of these techniques. For instance, Ruiz-Ramos (2016) analysed the application of several bias correction methods for improving crop impact projections in the Iberian Peninsula for 21st century. Gabaldon-Leal (2015) used bias corrected ENSEMBLES data to analyse summer crops in the southern Iberian Peninsula, focusing on the impacts of rising temperatures, and of higher frequency of extreme events on irrigated maize, and to evaluate some adaptation strategies.

Further research on bias correction methods include the work by Romero et al. (2011), presenting a new parametric trend-preserving bias correction method, and by Turco et al. (2017), introducing a new methodology for correcting and downscaling RCM data based on an analog technique.

Distillation of global and regional scenario results

VIA research communities are usually advised to consider an ensemble of model projections, or at least a selection of members spanning the ensemble uncertainty, in order to propagate the uncertainty arising from different greenhouse gases emission scenarios and (global and regional) climate models. However, the distillation of information out of the large amount of available data is a technical and ethical challenge (Hewitson et al., 2013). It is still not clear how to best proceed in order to select a subset of representative data (out from the large ensemble available) for a particular study (Cannon, 2015). Moreover, different datasets could provide inconsistent or even conflicting information making this process even harder. This is one of the key challenges considered in ongoing initiatives (e.g. the “distillation of climate information” work package within EURO-CORDEX), and further advances are expected in the coming years.

In order to illustrate this problem, we will see that the simple summary of ensemble members in percentile ranges shown in Figure 2 is misleading. Ensemble




Figure 3: Summer (JJA) temperature deltas (2021-2050 vs 1971-2000) averaged over continental Spain and the Balearic Islands. The 255 ensemble members considered in this work have been split into sub-ensembles considering only (a) GCMs, (b) SDMs and (c) RCMs. In each case, the lines at the bottom depict all sub-ensemble member deltas and are averaged at the joints as the plot progresses upward to the upper joint representing the sub-ensemble mean. Probability density functions are estimated for each sub-ensemble and 50% and 90% central range estimates are depicted as shades. See Fernández et al. (2017) for more information.

members are not independent and specific features of downscaling methods, and (lack of) ensemble design, are behind the different climate change ranges shown. As an example, we focus on summer temperature change (squared in Figure 2) and show individual results (Figure 3) for the three sources of future projections (GCMs, SDMs and RCMs). Individual delta change estimates at the bottom are coloured by the driving GCM, which is the largest source of uncertainty (spread). There are

GCMs, such as CanESM2, ranking in the upper end of the temperature change (Figure 3a), while others (e.g. CNRM-CM5, MPI-ESM) tend to project smaller changes. This trend is essentially preserved by downscaling methods (Figure 3b,c) with some particularities. In SDMs, the statistical method used (linear model -LM- or analogues -AN-) is a large source of spread, with analogue methods ranking clearly at the lower end due to their inability to extrapolate temperatures beyond the observed

range. On the other hand, the RCMs, which are more physically based, are computationally very expensive, and the available GCM population has been very unevenly sampled. In this example, RCMs have favoured two of the GCMs projecting the smallest delta changes (CNRM and MPI models). This fact, along with a genuine tendency by the RCMs to project smaller delta changes (potential added value) results in a narrower range and smaller mean temperature change than projected by GCMs.

Acknowledgements

We thank the modelling groups contributing to the CORDEX and CMIP initiatives, which provided data for this study through the ESGF infrastructure. The authors also want to thank, in particular, Petra Ramos (AEMET), Emma Gaitán (FIC), Jesús Asín (Univ. Zaragoza), Romu Romero (Univ. Islas Baleares) and Carmen Llasat (Univ. Barcelona) for providing a review of previous works.

References

Amengual, A., V. Homar, R. Romero, S. Alonso, and C. Ramis. 2011. "A Statistical Adjustment of Regional Climate Model Outputs to Local Scales: Application to Platja de Palma, Spain." *J. Climate* 25 (3): 939–57. doi:10.1175/JCLI-D-10-05024.1.

Bafaluy, D., A. Amengual, R. Romero, and V. Homar. 2014. "Present and Future Climate Resources for Various Types of Tourism in the Bay of Palma, Spain." *Regional Environmental Change* 14 (5): 1995–2006. doi:10.1007/s10113-013-0450-6.

Bedia, J., S. Herrera, A. Camia, J. M. Moreno, and J. M. Gutiérrez. 2014. "Forest Fire Danger Projections in the Mediterranean Using ENSEMBLES Regional Climate Change Scenarios." *Climatic Change* 122 (1–2): 185–199. doi:10.1007/s10584-013-1005-z.

Bedia, J., S. Herrera, D. San Martín, N. Koutsias, and J. M. Gutiérrez. 2013. "Robust Projections of Fire Weather Index in the Mediterranean Using Statistical Downscaling." *Climatic Change* 120 (1–2): 229–247. doi:10.1007/s10584-013-0787-3.

Cannon, A.J. 2015. "Selecting GCM Scenarios That Span the Range of Changes in a Multimodel Ensemble: Application to CMIP5 Climate Extremes Indices." *J. Climate* 28 (3): 1260–1267. doi:10.1175/JCLI-D-14-00636.1.

Casanueva, A., M. D. Frías, S. Herrera, D. San-Martín, K. Zaninovic, and J. M. Gutiérrez. 2014. "Statistical Downscaling of Climate Impact Indices: Testing the Direct Approach." *Climatic Change* 127 (3–4): 547–560. doi:10.1007/s10584-014-1270-5.

Casanueva, A., S. Kotlarski, S. Herrera, J. Fernández, J. M. Gutiérrez, F. Boberg, A. Colette, Christensen O.B., Goergen K., Jacob D., Keuler K., Nikulin G., Teichmann C.,

and Vautard R. "Daily Precipitation Statistics in a EURO-CORDEX RCM Ensemble: Added Value of Raw and Bias-Corrected High-Resolution Simulations." *Clim. Dyn.* 47 (3–4): 719–37. doi:10.1007/s00382-015-2865-x.

Déqué, M., S. Somot, E. Sanchez-Gomez, C. M. Goodess, D. Jacob, G. Lenderink, and O. B. Christensen. 2012. "The Spread amongst ENSEMBLES Regional Scenarios: Regional Climate Models, Driving General Circulation Models and Interannual Variability." *Clim. Dyn.* 38 (5–6): 951–964. doi:10.1007/s00382-011-1053-x.

Domínguez, M., R. Romera, E. Sánchez, L. Fita, J. Fernández, P. Jiménez-Guerrero, J. P. Montávez, W. D. Cabos, G. Liguori, M. A. Gaertner. 2013. "Present-climate precipitation and temperature extremes over Spain from a set of high resolution RCMs". *Clim. Res.*, 58(2):149–164.

Dosio, A., P. Paruolo, and R. Rojas. 2012. "Bias Correction of the ENSEMBLES High Resolution Climate Change Projections for Use by Impact Models: Analysis of the Climate Change Signal." *J. Geophys. Res.* 117 (D17). doi:10.1029/2012JD017968

Escenarios-PNACC (2012) data: [1] http://www.aemet.es/es/serviciosclimaticos/cambio_climat/datos_mensuales/ayuda; [2] <http://www.meteo.unican.es/escenarios-pnacc>

Esteve-Selma, M. A., Martínez-Fernández, J., Hernández-García, I., Montávez, J. P., López-Hernández, J. J., & Calvo, J. F. (2012). Potential effects of climatic change on the distribution of *Tetraclinis articulata*, an endemic tree from arid Mediterranean ecosystems. *Climatic change*, 113(3–4), 663–678.

Fernández, J. , L. Fita, M. García-Díez, J. P. Montávez, P. Jiménez-Guerrero, M. Domínguez, R. Romera, N. López de la Franca, E. Sánchez, G. Liguori, W. D. Cabos, M. A. Gaertner, 2012. Escenarios-PNACC 2012: Resultados de regionalización dinámica. En "Cambio climático: Extremos e Impactos". Publicaciones de la AEC. Serie A, n8 (in Spanish).

Fernández, J. , M. D. Frías, W. D. Cabos, A. S. Cofiño, M. Domínguez, L. Fita, M. A. Gaertner, M. García-Díez, J. M. Gutiérrez, P. Jiménez-Guerrero, G. Liguori, J. P. Montávez, R. Romera, E. Sánchez., 2017. Consistency of climate change projections from multiple global and regional model intercomparison projects. Submitted to *Climatic Change*.

Frías, M.D., R. Mínguez, J.M. Gutiérrez, and F.J. Méndez. (2012). Future Regional Projections of Extreme Temperatures in Europe: A Nonstationary Seasonal Approach. *Climatic Change* 113 (2): 371–392. doi:10.1007/s10584-011-0351-y.

- Gabaldón-Leal, C., Lorite, I. J., Minguez Tudela, M. I., Lizaso Oñate, J. I., Dosio, A., Sánchez, E., Ruiz Ramos, M. (2015). Strategies for adapting maize to climate change and extreme temperatures in Andalusia, Spain. *Climate Research*, 65, 159-173.
- Gómez, G., Cabos, W.D., Liguori, G., Sein, D., Lozano-Galeana, S., Fita, L., Fernández, J., Magariño, M.E., Jiménez-Guerrero, P., Montávez, J.P. and Domínguez, M. 2016. Characterization of the wind speed variability and future change in the Iberian Peninsula and the Balearic Islands. *Wind Energy*, 19(7), 1223-1237.
- Gómez-Navarro, J. J., Montávez, J. P., Jimenez-Guerrero, P., Jerez, S., Garcia-Valero, J. A., & González-Rouco, J. F. (2010). Warming patterns in regional climate change projections over the Iberian Peninsula. *Meteorologische Zeitschrift*, 19(3), 275-285.
- Gonçalves, M., A. Barrera-Escoda, D. Guerreiro, J. M. Baldasano, and J. Cunillera. 2014. "Seasonal to Yearly Assessment of Temperature and Precipitation Trends in the North Western Mediterranean Basin by Dynamical Downscaling of Climate Scenarios at High Resolution (1971–2050)." *Climatic Change* 122 (1–2): 243–56. doi:10.1007/s10584-013-0994-y.
- Gutiérrez, J.M. , J. Ribalaygua, C. Llasat, R. Romero, J. Abaurrea, E. Rodríguez. 2012. Escenarios-PNACC 2012: Descripción y Análisis de los Resultados de Regionalización Estadística. En "Cambio climático: Extremos e Impactos". Publicaciones de la AEC. Serie A, n8, 125-134. ISBN: 978-84-695-4331-3 (in Spanish).
- Gutiérrez, J. M., D. San-Martín, S. Brands, R. Manzanás, and S. Herrera. 2013. "Reassessing Statistical Downscaling Techniques for Their Robust Application under Climate Change Conditions." *J. Climate* 26 (1): 171–188. doi:10.1175/JCLI-D-11-00687.1.
- Herrera, S., J. M. Gutiérrez, R. Ancell, M. R. Pons, M. D. Frías, and J. Fernández. 2012. "Development and Analysis of a 50-Year High-Resolution Daily Gridded Precipitation Dataset over Spain (Spain02)." *Int. J. Climatol.* 32 (1): 74–85. doi:10.1002/joc.2256.
- Hewitson, B. C., J. Daron, R. G. Crane, M. F. Zermoglio, and C. Jack. 2013. "Interrogating Empirical-Statistical Downscaling." *Climatic Change* 122 (4): 539–54. doi:10.1007/s10584-013-1021-z.
- Hewitson, Bruce, Katinka Waagsaether, Jan Wohland, Kate Kloppers, and Teizeen Kara. 2017. "Climate Information Websites: An Evolving Landscape." *Wiley Interdisciplinary Reviews: Climate Change*. doi:10.1002/wcc.470.
- Jacob, D., J. Petersen, B. Eggert, A. Alias, O. B. Christensen, L. M. Bouwer, A. Braun, et al. 2014. "EURO-CORDEX: New High-Resolution Climate Change Projections for European Impact Research." *Regional Environmental Change* 14 (2): 563–78. doi:10.1007/s10113-013-0499-2.
- Jerez, S., J. Montávez, J. Gómez-Navarro, P. Jiménez, P. Jiménez-Guerrero, R. Lorente, and J. F. Gonzalez-Rouco. 2012. "The role of the land-surface model for climate change projections over the Iberian Peninsula". *J. Geophys. Res. : Atmospheres*, 117(D1).
- Jerez, S., J. P. Montávez, J. J. Gómez-Navarro, R. Lorente-Plazas, J. A. Garcia-Valero, P. Jimenez-Guerrero. 2013. "A multi-physics ensemble of regional climate change projections over the Iberian Peninsula". *Clim. Dyn.*, 41(7-8):1749–1768.
- Jerez, S., Tobin, I., Vautard, R., Montávez, J.P., López-Romero, J.M., Thais, F., Bartok, B., Christensen, O.B., Colette, A., Déqué, M. and Nikulin, G. 2015. "The impact of climate change on photovoltaic power generation in Europe". *Nature communications*, 6, 10014.
- Jiménez-Guerrero, P., J. Montávez, M. Domínguez, R. Romera, L. Fita, J. Fernández, W. Cabos, G. Liguori, and M. Gaertner, 2013. Mean fields and interannual variability in rcm simulations over Spain: the escena project. *Climate Research*, 57(3):201–220.
- López-Franca, N., E. Sánchez, T. Losada, M. Domínguez, R. Romera, M. Á. Gaertner. 2015. "Markovian Characteristics of Dry Spells over the Iberian Peninsula under Present and Future Conditions using ESCENA Ensemble of Regional Climate Models." *Clim. Dyn.* 45 (3–4): 661–77. doi:10.1007/s00382-014-2280-8.
- Maraun, D., M. Widmann, J. M. Gutiérrez, S. Kotlarski, R. E. Chandler, E. Hertig, J. Wibig, R. Huth, R.A.I. Wilcke. 2015. "VALUE: A Framework to Validate Downscaling Approaches for Climate Change Studies." *Earth's Future* 3 (1): 2014EF000259. doi:10.1002/2014EF000259.
- Monjo, R., Gaitán, E., Pórtolles, J., Ribalaygua, J., Torres, L. 2016. "Changes in extreme precipitation over Spain using statistical downscaling of CMIP5 projections". *Int. J. Climatol.*, 36: 757–769. doi:10.1002/joc.4380
- Osca, J., R. Romero, and S. Alonso. 2013. "Precipitation Projections for Spain by Means of a Weather Typing Statistical Method." *Global and Planetary Change* 109 (October): 46–63. doi:10.1016/j.gloplacha.2013.08.001.
- Pérez, J. C., J. P. Díaz, A. González, J. Expósito, F. Rivera-López, and D. Taima. 2014. "Evaluation of WRF Parameterizations for Dynamical Downscaling in the Canary Islands." *J. Climate* 27 (14): 5611–31. doi:10.1175/JCLI-D-13-00458.1.

- Pons, M. R., S. Herrera, and J. M. Gutiérrez. 2016. "Future Trends of Snowfall Days in Northern Spain from ENSEMBLES Regional Climate Projections." *Clim. Dyn.* 46 (11–12): 3645–55. doi:10.1007/s00382-015-2793-9.
- Ramos, P., Petisco, E., Martín, J. M., Rodríguez, E. 2013. "Downscaled climate change projections over Spain: application to water resources". *Int. J. Water Res. Devel.*, 29(2), 201-218
- Resco, P., A. Iglesias, I. Bardají, and V. Sotés. 2016. "Exploring Adaptation Choices for Grapevine Regions in Spain." *Regional Environmental Change* 16 (4): 979–93. doi:10.1007/s10113-015-0811-4.
- Ribalaygua, J., M.R. Pino, J. Pórtolles, E. Roldán, E. Gaitán, D. Chinarro, L. Torres. 2013. "Climate Change Scenarios for Temperature and Precipitation in Aragón (Spain)." *Science of The Total Environment* 463 (October): 1015–30. doi:10.1016/j.scitotenv.2013.06.089.
- Romera, R., M.Á. Gaertner, E. Sánchez, M. Domínguez, J.J. González-Alemán, M.M. Miglietta. 2017. "Climate change projections of medicanes with a large multi-model ensemble of regional climate models". *Glob.Planet.Change*, 151: 134-143, doi: 10.1016/j.gloplacha.2016.10.008.
- Romero, R., and K. Emanuel. 2017. "Climate Change and Hurricane-Like Extratropical Cyclones: Projections for North Atlantic Polar Lows and Medicanes Based on CMIP5 Models". *J. Climate*. 30(1): 279-299.
- Ruiz-Ramos, M., A. Rodríguez, A. Dosio, C. M. Goodess, C. Harpham, M. I. Mínguez, and E. Sánchez. 2016. "Comparing Correction Methods of RCM Outputs for Improving Crop Impact Projections in the Iberian Peninsula for 21st Century." *Climatic Change* 134 (1–2): 283–97. doi:10.1007/s10584-015-1518-8.
- Ruti, P., P. M. Ruti, S. Somot, F. Giorgi, C. Dubois, E. Flaounas, A. Obermann, A. Dell'Aquila, G. Pisacane, A. Harzallah, E. Lombardi, B. Ahrens, N. Akhtar, A. Alias, T. Arsouze, R. Aznar, S. Bastin, J. Bartholy, K. Béranger, J. Beuvier, S. Bouffies-Cloch  , J. Brauch, W. Cabos, S. Calmanti, J.-C. Calvet, A. Carillo, D. Conte, E. Coppola, V. Djurdjevic, P. Drobinski, A. Elizalde-Arellano, M. Gaertner, P. Gal  n, C. Gallardo, S. Gualdi, M. Goncalves, O. Jorba, G. Jord  , B. L'Heveder, C. Lebeau-pin-Brossier, L. Li, G. Liguori, P. Lionello, D. Maci  s, P. Nabat, B.   nol, B. Raikovic, K. Ramage, F. Sevault, G. Sannino, M. V. Struglia, A. Sanna, C. Torma, V. Vervatis., 2016. "MED-CORDEX initiative for Mediterranean Climate studies". *Bull. Amer. Met. Soc.*, 2016, 1187-1208.
- S  nchez E., and G. Miguez-Macho. 2010. Regional climate projections over the Iberian Peninsula: climate change scenarios modeling. CLIVAR-Spain Regional climate change assessment report: Climate in Spain: Past, Present and Future (Editors: P  rez F. Fiz and Boscolo Roberta) pp. 69-80. Available at <http://www.clivar.org/documents/spain-2010>
- San-Mart  n, D., R. Manzan  s, S. Brands, S. Herrera, and J. M. Guti  rrez. 2017. "Reassessing Model Uncertainty for Regional Projections of Precipitation with an Ensemble of Statistical Downscaling Methods." *J. Climate* 30 (1): 203–23. doi:10.1175/JCLI-D-16-0366.1.
- Turco, M., M. C. Llasat, S. Herrera, and J. M. Guti  rrez. 2017. "Bias Correction and Downscaling of Future RCM Precipitation Projections Using a MOS-Analog Technique." *J. Geophys. Res.: Atmospheres* 122 (5): 2631–48. doi:10.1002/2016JD025724.
- Turco, M., P. Quintana-Segui, M. C. Llasat, S. Herrera, and J. M. Guti  rrez. 2011. "Testing MOS Precipitation Downscaling for ENSEMBLES Regional Climate Models over Spain." *J. Geophys. Res.* 116 (D18). doi:10.1029/2011JD016166.
- Turco, M., A. Sanna, S. Herrera, M.C. Llasat, J.M. Guti  rrez. 2015. "Evaluation of the ENSEMBLES Transient RCM Simulations Over Spain: Present Climate Performance and Future Projections." In *Engineering Geology for Society and Territory - Volume 1*, edited by G. Lollino, et al. 199–203. Springer International Publishing.